

### Acute Coronary Syndrome (AMI/Chest Pain)

Admitting Physician: \_\_\_\_\_ Date: \_\_\_\_\_ Time: \_\_\_\_\_

**Admit to Inpatient:** [ ] Med/Surg [ ] Telemetry [ ] ICU  
**Place in Observation Services:** [ ] Med/Surg [ ] Telemetry [ ] Other: \_\_\_\_\_  
**Diagnosis:** [ ] Acute Coronary Syndrome [ ] Chest Pain [ ] Acute MI [ ] Other: \_\_\_\_\_  
**Condition:** [ ] Stable [ ] Guarded [ ] Critical [ ] Good [ ] Fair [ ] Poor  
**Consult:** [ ] Cardiology: \_\_\_\_\_  
**Allergies:** [ ] Cardiovascular Surgery: \_\_\_\_\_  
 [ ] Other: \_\_\_\_\_

**Code Status:** [ ] Full [ ] DNR

**Vital Signs:** [ ] Per unit protocol [ ] Every shift [ ] every \_\_\_\_\_ hours  
 [ ] Other: \_\_\_\_\_

**Activity:** [ ] Bed rest [ ] Up in chair [ ] Bedside commode [ ] Ambulate ad lib [ ] Bathroom privileges

**Nursing:** [ ] Daily weight [ ] Intake & Output [ ] Foley to drainage [ ] Sequential Compression Device (SCD)  
 [ ] Pulse oximeter [ ] every shift [ ] twice a day [ ] Continuous monitoring  
 [ ] Glucose checks AC and at bedtime or every \_\_\_\_\_ hours  
 [ ] Other: \_\_\_\_\_

**Diet:** [ ] Regular [ ] Mechanical soft [ ] Restrict fluid \_\_\_\_\_ [ ] Sodium restricted \_\_\_\_\_ mg  
 [ ] Cardiac/Heart Healthy [ ] Diabetic (specify) \_\_\_\_\_ calorie [ ] Renal/Kidney  
 [ ] Other: \_\_\_\_\_

**Fluids:** [ ] Intravenous: \_\_\_\_\_ @ \_\_\_\_\_ ml/hr  
 [ ] Saline lock [ ] Other: \_\_\_\_\_

**Oxygen:** [ ] Nasal Cannula \_\_\_\_\_ L/min, adjust to keep O2 sat greater than 90%.  
 [ ] Venti Mask \_\_\_\_\_ % FIO2 [ ] 100% NRB  
 [ ] Other: \_\_\_\_\_

**Labs:** [ ] Troponin (STAT, repeat 90 min, and 12 hours) [ ] CPK [ ] Myoglobin  
 [ ] BMP [ ] CMP [ ] BNP [ ] Magnesium [ ] Stool guaiac  
 [ ] Fasting lipid profile [ ] TSH [ ] UA [ ] CBC [ ] ABG  
 [ ] D-dimer [ ] Fibrinogen [ ] High Sensitivity C-Reactive Protein  
 [ ] Other labs: \_\_\_\_\_

**Protocols (If available): All protocol orders must be placed in chart**

[ ] Weight Based Heparin Dosing Protocol (when patient placed on protocol, all associated labs and monitorin is included)  
 [ ] Insulin Infusion Protocol (ICU patients only)

**Studies: (If LVEF less than 40%, Implement CHF Protocol**

[ ] Cardiac Stress Test: (after negative enzymes x3)  
 [ ] CXR: [ ] Portable [ ] PA/Lateral [ ] Exercise treadmill  
 [ ] EKG STAT, repeat in 12 hours and in AM [ ] Stress echocardiogram  
 [ ] STAT EKG PRN with chest pain or palpitations [ ] Adenosine treadmill  
 [ ] Echocardiogram Dr. \_\_\_\_\_ to read [ ] Persantin Thallium  
 [ ] Nuclear Cardiac Scan (assess wall motion, EF) [ ] Dobutamine Stress Test  
 [ ] Other: \_\_\_\_\_ [ ] Treadmill with Cardiolyte

NURSE NOTED	DATE	TIME	DATE / TIME	PHYSICIAN SIGNATURE OR AUTHENTICATION
24 HR. CHART CHECK BY NURSE	DATE	TIME		



Account Number: \_\_\_\_\_ MR Number: \_\_\_\_\_  
 Patient Name: \_\_\_\_\_  
 Admit Date: \_\_\_\_\_



**Salt Lake Regional**  
**MEDICAL CENTER**

1050 East South Temple - Salt Lake City - Utah 84102  
 (801) 350-4111

DOB	Age	Sex	HT	WT	RM-BD	PT	SVC	FC
Allergies: _____								
Attending Physician Name: _____								

**Medications: Check the appropriate box.**

- Aspirin 325 mg (81 mg chewable tablets x 4) PO now, then  81 mg PO daily  162 mg PO daily
- Clopidogrel (Plavix) \_\_\_\_\_ mg PO x 1 now, then 75 mg PO once daily
- Morphine Sulfate 2 mg IV every 10 minutes PRN severe pain (max 8 mg/hr)
- Nitroglycerin 0.4mg SL every 5 minutes as needed for chest pain (Max of 3 tablets in 15 minutes)
- Nitroglycerin infusion 50 mg/250 ml D5W at 200 micrograms/ml
- Titrate to control symptoms in  5 microgram/min  10 microgram/min steps,  
up to a maximum of \_\_\_\_\_ microgram/kg per min; maintain SBP more than 100mm
- Nitroglycerin paste  1 inch  2 inch topically to chest wall every 6 hours. Remove at bedtime.
- Isosorbide dinitrate (Isordil)  5 mg PO TID  10 mg PO TID  20 mg PO TID  30 mg PO TID  40 mg PO TID  60 mg PO TID
- Enoxaparin (Lovenox) 40 mg subcutaneous every 24 hrs
- Amiodarone 150mg IV bolus over 10 minutes. Start infusion at 1 mg/min for 6 hrs, then 0.5 mg/min for 18 hrs
- Eptifibatid (Integrilin) 180 microgram/kg IV push, then 2 microgram/kg per minute for  48 hrs  72 hrs
- Metoprolol (Lopressor) 5 mg IV every 5 minutes x 3 doses
- Metoprolol (Lopressor) 25 mg PO BID. Start in AM
- Lisinopril (Zestril, Prinivil) 5 mg PO now, then 10 mg daily (start in AM).
- If allergic or intolerant to ACEI, use Losartan (Cozaar) 25mg PO daily.
- Reteplase (Retavase) 10 units IV bolus over 2 minutes. Repeat bolus in 30 minutes
- Simvastatin (Zocor) 40 mg PO, every night before bedtime
- Lorazepam (Ativan)  0.5mg PO every 6 PRN  1 mg PO every 6 PRN hrs as needed for anxiety
- Famotidine (Pepcid) 20 mg  IV twice daily  PO twice daily
- Pantoprazole (Protonix) 40mg IV once daily
- Nicotine \_\_\_\_\_ mg topical patch apply daily
- Percocet 5/325 mg PO every 4 hrs PRN for moderate pain
- Morphine Sulfate 2 mg IV every 4 hr PRN severe pain (unless patient is allergic to codeine or morphine)
- Temazepam (Restoril) 15 mg PO bedtime PRN sleep. May repeat dose once in one hour if no results.
- Ondansetron (Zofran) 4 mg IV every 24 hrs PRN for nausea and vomiting



**Standard Medications: (all orders below will be implemented unless crossed out)**


- Acetaminophen (Tylenol) 650 mg PO every 4 hr PRN mild pain or temp greater than 101F (not to exceed 4 grams per 24 hrs)
- Docusate sodium (Colace) 100 mg PO every HS
- MOM 30 ml PO PRN constipation
- Maalox 30 ml PO PRN heartburn

**Additional Meds:**

- \_\_\_\_\_
- \_\_\_\_\_
- \_\_\_\_\_
- \_\_\_\_\_
- \_\_\_\_\_

NURSE NOTED	DATE	TIME	DATE / TIME	PHYSICIAN SIGNATURE OR AUTHENTICATION
24 HR. CHART CHECK BY NURSE	DATE	TIME		

 	Account Number:	MR Number:
Patient Name:		
Admit Date:		

 <p><b>Salt Lake Regional</b> MEDICAL CENTER</p> <p>1050 East South Temple - Salt Lake City - Utah 84102 (801) 350-4111</p>	DOB	Age	Sex	HT	WT	RM-BD	PT	SVC	FC
	Allergies:								
	Attending Physician Name:								